

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number
WO 03/028466 A2

(51) International Patent Classification⁷: **A01N 43/90**,
25/32 // (A01N 43/90, 47:36, 43:824, 35:06)

(21) International Application Number: PCT/EP02/10829

(22) International Filing Date:
26 September 2002 (26.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1781/01 27 September 2001 (27.09.2001) CH

(71) Applicant (for all designated States except US): **SYNGENTA PARTICIPATIONS AG** [CH/CH]; Schwarzwaldallee 215, 4058 Basel (CH).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **HOFER, Urs** [CH/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, 4058 Basel (CH).

(74) Agent: **BASTIAN, Werner**; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **HERBICIDAL COMPOSITION**



(I)


(57) Abstract: Herbicidal composition comprising: a) a herbicidally effective amount of a compound of formula (I), wherein the substituents are as defined in claim 1; and b) an amount, effective for herbicide synergy, of mesosulfuron, mesotrione or flufenacet.

Herbicidal composition

The present invention relates to new herbicidal compositions for controlling grasses and weeds in crops of useful plants, especially in crops of maize and cereals, which compositions comprise a tetrahydropyrazolodione herbicide known from, for example, WO 99/47525, and a co-herbicide.

The present invention relates to a herbicidal composition that, in addition to comprising customary inert formulation adjuvants, such as carriers, solvents and wetting agents, comprises as active ingredient a mixture of


a) a herbicidally effective amount of a compound of formula I



wherein

R₁ and R₃ are, each independently of the other, halogen, nitro, cyano, C₁-C₄alkyl, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkyl, C₂-C₆haloalkenyl, C₃-C₆cycloalkyl, halo-substituted C₃-C₆cycloalkyl, C₂-C₆alkoxyalkyl, C₂-C₆alkylthioalkyl, hydroxy, mercapto, C₁-C₆alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkynyloxy, carbonyl, carboxyl, C₁-C₄alkylcarbonyl, C₁-C₄hydroxyalkyl, C₁-C₄alkoxycarbonyl, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, amino, C₁-C₄alkylamino or di(C₁-C₄alkyl)amino;

R₄ and R₅ together are a group



wherein R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈ and R₂₉ are, each independently of the others, hydrogen, halogen, C₁-C₄alkyl or C₁-C₄haloalkyl, it being possible for an alkylene ring, which together with the carbon atoms of the groups Z₁, Z₂ or Z₃ contains from 2 to 6 carbon atoms and which may be interrupted by

oxygen, to be either fused or spiro-bound to the carbon atoms of the groups Z_1 , Z_2 or Z_3 , or that alkylene ring bridges at least one ring atom of the groups Z_1 , Z_2 or Z_3 ;

G is hydrogen, $-C(X_1)-R_{30}$, $-C(X_2)-X_3-R_{31}$, $-C(X_4)-N(R_{32})-R_{33}$, $-SO_2-R_{34}$, an alkali metal, alkaline earth metal, sulfonium or ammonium cation, or $-P(X_5)(R_{35})-R_{36}$ or $-CH_2-X_6-R_{37}$;

X_1 , X_2 , X_3 , X_4 , X_5 and X_6 are, each independently of the others, oxygen or sulfur;

R_{30} , R_{31} , R_{32} and R_{33} are, each independently of the others, hydrogen, C_1 - C_{10} alkyl, C_1 - C_{10} haloalkyl, C_1 - C_{10} cyanoalkyl, C_1 - C_{10} nitroalkyl, C_1 - C_{10} aminoalkyl, C_1 - C_5 alkylamino- C_1 - C_5 alkyl, C_2 - C_8 dialkylamino- C_1 - C_5 alkyl, C_3 - C_7 cycloalkyl- C_1 - C_5 alkyl, C_2 - C_{10} alkoxy-alkyl, C_4 - C_{10} alkenyloxy-alkyl, C_4 - C_{10} alkynyloxy-alkyl, C_2 - C_{10} alkylthio-alkyl, C_1 - C_5 alkylsulfoxyl- C_1 - C_5 alkyl, C_1 - C_5 alkylsulfonyl- C_1 - C_5 alkyl, C_2 - C_8 alkylideneamino-oxy- C_1 - C_5 alkyl, C_1 - C_5 alkylcarbonyl- C_1 - C_5 alkyl, C_1 - C_5 alkoxycarbonyl- C_1 - C_5 alkyl, C_1 - C_5 amino-carbonyl- C_1 - C_5 alkyl, C_2 - C_8 dialkylamino-carbonyl- C_1 - C_5 alkyl, C_1 - C_5 alkylcarbonylamino- C_1 - C_5 alkyl, C_2 - C_5 alkylcarbonyl- $(C_1$ - C_5 -alkyl)-aminoalkyl, C_3 - C_6 trialkylsilyl- C_1 - C_5 alkyl, phenyl- C_1 - C_5 alkyl, heteroaryl- C_1 - C_5 alkyl, phenoxy- C_1 - C_5 alkyl, heteroaryloxy- C_1 - C_5 alkyl, C_2 - C_5 alkenyl, C_2 - C_5 haloalkenyl, C_3 - C_8 cycloalkyl, phenyl, or C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted phenyl or heteroaryl or heteroaryl-amino, C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted heteroaryl-amino, diheteroaryl-amino, C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted diheteroaryl-amino, phenylamino, C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted phenylamino, diphenylamino, C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted diphenyl-amino, C_3 - C_7 cycloalkylamino, C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted C_3 - C_7 cycloalkylamino, di- C_3 - C_7 cycloalkylamino, C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted di- C_3 - C_7 cycloalkylamino, C_3 - C_7 cycloalkoxy or C_1 - C_3 alkyl-, C_1 - C_3 haloalkyl-, C_1 - C_3 alkoxy-, C_1 - C_3 haloalkoxy-, halo-, cyano- or nitro-substituted C_3 - C_7 cycloalkoxy;

R_{34} , R_{35} and R_{36} are, each independently of the others, hydrogen, C_1 - C_{10} alkyl, C_1 - C_{10} haloalkyl, C_1 - C_{10} cyanoalkyl, C_1 - C_{10} nitroalkyl, C_1 - C_{10} aminoalkyl, C_1 - C_5 alkylamino- C_1 - C_5 alkyl, C_2 - C_8 dialkylamino- C_1 - C_5 alkyl, C_3 - C_7 cycloalkyl- C_1 - C_5 alkyl, C_2 - C_{10} alkoxy-alkyl, C_4 - C_{10} alkenyloxy-alkyl, C_4 - C_{10} alkynyloxy-alkyl, C_2 - C_{10} alkylthio-alkyl, C_1 - C_5 alkylsulfoxyl- C_1 - C_5 alkyl, C_1 - C_5 alkylsulfonyl- C_1 - C_5 alkyl, C_2 - C_8 alkylideneamino-oxy- C_1 - C_5 alkyl, C_1 - C_5 alkylcarbonyl- C_1 - C_5 alkyl, C_1 - C_5 alkoxycarbonyl- C_1 - C_5 alkyl, C_1 - C_5 amino-carbonyl- C_1 - C_5 alkyl, C_2 - C_8 dialkylamino-carbonyl- C_1 - C_5 alkyl, C_1 - C_5 alkylcarbonylamino- C_1 - C_5 alkyl, C_2 - C_5 alkylcarbonyl-

(C₁-C₅alkyl)-aminoalkyl, C₃-C₆trialkylsilyl-C₁-C₅alkyl, phenyl-C₁-C₅alkyl, heteroaryl-C₁-C₅alkyl, phenoxy-C₁-C₅alkyl, heteroaryloxy-C₁-C₅alkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, phenyl, or C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl or heteroaryl or heteroarylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted heteroarylamino, diheteroaryl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diheteroarylamino, phenylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenylamino, diphenylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diphenylamino, C₃-C₇cycloalkylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkylamino, di-C₃-C₇cycloalkylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted di-C₃-C₇cycloalkylamino, C₃-C₇cycloalkoxy, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀haloalkoxy, C₁-C₅alkylamino, C₂-C₈dialkylamino and benzyloxy or phenoxy, it being possible for the benzyl and phenyl groups themselves to be substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano, formyl, acetyl, propionyl, carboxyl, C₁-C₅alkoxy-carbonyl, methylthio, ethylthio or by nitro;

and

R₃₇ is C₁-C₁₀alkyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino-C₁-C₅alkyl, C₂-C₈dialkylamino-C₁-C₅alkyl, C₃-C₇cycloalkyl-C₁-C₅alkyl, C₂-C₁₀alkoxy-alkyl, C₄-C₁₀alkenyloxy-alkyl, C₄-C₁₀alkynyloxy-alkyl, C₂-C₁₀alkylthio-alkyl, C₁-C₅alkylsulfoxyl-C₁-C₅alkyl, C₁-C₅alkylsulfonyl-C₁-C₅alkyl, C₂-C₈alkylideneamino-oxy-C₁-C₅alkyl, C₁-C₅alkylcarbonyl-C₁-C₅alkyl, C₁-C₅alkoxycarbonyl-C₁-C₅alkyl, C₁-C₅amino-carbonyl-C₁-C₅alkyl, C₂-C₈dialkylamino-carbonyl-C₁-C₅alkyl, C₁-C₅alkylcarbonylamino-C₁-C₅alkyl, C₂-C₅alkylcarbonyl-(C₁-C₅alkyl)-aminoalkyl, C₃-C₆trialkylsilyl-C₁-C₅alkyl, phenyl-C₁-C₅alkyl, heteroaryl-C₁-C₅alkyl, phenoxy-C₁-C₅alkyl, heteroaryloxy-C₁-C₅alkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, phenyl, or C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl or heteroaryl, or heteroarylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted heteroarylamino, diheteroarylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diheteroarylamino, phenylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl-

amino, diphenylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diphenylamino, C₃-C₇cycloalkylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkylamino, di-C₃-C₇cycloalkylamino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted di-C₃-C₇cycloalkylamino, C₃-C₇cycloalkoxy, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkoxy or C₁-C₁₀alkylcarbonyl; and salts and diastereoisomers of the compounds of formula I, with the proviso that R₁ and R₃ are not simultaneously methyl; and

b) an amount, effective for herbicide synergy, of at least one herbicide selected from mesosulfuron, mesotrione and flufenacet.

In the above definitions, halogen is to be understood as fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine. The alkyl groups occurring in the substituent definitions are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl or tert-butyl, and the pentyl and hexyl isomers. Suitable cycloalkyl substituents contain from 3 to 6 carbon atoms and are, for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl. They may be substituted one or more times by halogen, preferably fluorine, chlorine or bromine. Alkenyl is to be understood as, for example, vinyl, allyl, methallyl, 1-methylvinyl or but-2-en-1-yl. Alkynyl is, for example, ethynyl, propargyl, but-2-yn-1-yl, 2-methylbutyn-2-yl or but-3-yn-2-yl. Haloalkyl groups preferably have a chain length of from 1 to 4 carbon atoms. Haloalkyl is, for example, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 2-fluoroethyl, 2-chloroethyl, pentafluoroethyl, 1,1-difluoro-2,2,2-trichloroethyl, 2,2,3,3-tetrafluoroethyl or 2,2,2-trichloroethyl, preferably trichloromethyl, difluorochloromethyl, difluoromethyl, trifluoromethyl or dichlorofluoromethyl. Suitable haloalkenyl radicals include alkenyl groups substituted one or more times by halogen, halogen being fluorine, chlorine, bromine or iodine and especially fluorine or chlorine, for example 2,2-difluoro-1-methylvinyl, 3-fluoropropenyl, 3-chloropropenyl, 3-bromopropenyl, 2,3,3-trifluoropropenyl, 2,3,3-trichloropropenyl and 4,4,4-trifluorobut-2-en-1-yl. Among the C₂-C₆alkenyl groups substituted once, twice or three times by halogen preference is given to those having a chain length of from 3 to 5 carbon atoms. Alkoxy groups preferably have a chain length of from 1 to 6 carbon atoms. Alkoxy is, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy or tert-butoxy, or a pentyloxy or hexyloxy isomer, preferably methoxy or ethoxy. Alkylcarbonyl is preferably

acetyl or propionyl. Alkoxy-carbonyl is, for example, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl or tert-butoxycarbonyl, preferably methoxycarbonyl or ethoxycarbonyl. Alkylthio groups preferably have a chain length of from 1 to 4 carbon atoms. Alkylthio is, for example, methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, isobutylthio, sec-butylthio or tert-butylthio, preferably methylthio or ethylthio. Alkylsulfinyl is, for example, methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl, n-butylsulfinyl, isobutylsulfinyl, sec-butylsulfinyl or tert-butylsulfinyl, preferably methylsulfinyl or ethylsulfinyl. Alkylsulfonyl is, for example, methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, isobutylsulfonyl, sec-butylsulfonyl or tert-butylsulfonyl, preferably methylsulfonyl or ethylsulfonyl. Alkylamino is, for example, methylamino, ethylamino, n-propylamino, isopropylamino or a butylamine isomer. Dialkylamino is, for example, dimethylamino, methylethylamino, diethylamino, n-propylmethylamino, dibutylamino or diisopropylamino. Alkoxyalkyl groups preferably have from 2 to 6 carbon atoms. Alkoxyalkyl is, for example, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, n-propoxymethyl, n-propoxyethyl, isopropoxymethyl or isopropoxyethyl. Alkylthioalkyl is, for example, methylthiomethyl, methylthioethyl, ethylthiomethyl, ethylthioethyl, n-propylthiomethyl, n-propylthioethyl, isopropylthiomethyl, isopropylthioethyl, butylthiomethyl, butylthioethyl or butylthiobutyl. Phenyl may be in substituted form, in which case the substituents may be in the ortho-, meta- and/or para-position. Preferred positions for the substituents are the ortho- and para-positions to the ring attachment point. Heteroaryl groups are usually aromatic heterocycles that contain preferably from 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur. Examples of suitable heterocycles and heteroaromatic compounds are: pyrrolidine, piperidine, pyran, dioxane, azetidine, oxetane, pyridine, pyrimidine, triazine, thiazole, thiadiazole, imidazole, oxazole, isoxazole and pyrazine, furan, morpholine, piperazine, pyrazole, benzoxazole, benzothiazole, quinoxaline and quinoline. Those heterocycles and heteroaromatic compounds may be further substituted, for example by halogen, alkyl, alkoxy, haloalkyl, haloalkoxy, nitro, cyano, thioalkyl, alkylamino or by phenyl. The C₂-C₁₀-alkenyl and -alkynyl groups R₃₄ may be mono- or poly-unsaturated. They contain preferably from 2 to 12 carbon atoms, especially from 2 to 6 carbon atoms.


Alkali metal, alkaline earth metal or ammonium cations for the substituent G are, for example, the cations of sodium, potassium, magnesium, calcium and ammonium. Preferred

- 6 -


sulfonium cations are especially trialkylsulfonium cations wherein the alkyl radicals each contain preferably from 1 to 4 carbon atoms.

The left-hand free valence of the groups Z_1 , Z_2 and Z_3 is bound to the 1-position and the right-hand free valence to the 2-position of the pyrazoline ring.

Compounds of formula I wherein it is possible for an alkylene ring, which together with the carbon atoms of the groups Z_1 , Z_2 and Z_3 contains from 2 to 6 carbon atoms, to be fused or spiro-bound to the groups Z_1 , Z_2 and Z_3 have, for example, the following structure:




(spiro-bound) or



(fused).

Compounds of formula I wherein in the groups Z_1 , Z_2 or Z_3 , an alkylene ring bridges at least one ring atom of the groups Z_1 , Z_2 or Z_3 , have, for example, the following structure:



(bridged).

Flufenacet is known from The Pesticide Manual, 12th Edition (BCPC) 2000, Entry No. 362; mesotrione is known from The Pesticide Manual, 12th Edition (BCPC) 2000, Entry No. 500; mesosulfuron is described, for example, in WO 00/3591 and WO 01/24633.

In herbicides of formula I that are preferred for the compositions in accordance with the invention, R_1 and R_3 are, each independently of the other, ethyl, haloethyl, ethynyl, C_1 - C_2 -alkoxy or C_1 - C_2 -haloalkoxy.

Preference is given also to those compositions in accordance with the invention wherein, in the herbicides of formula I, R_4 and R_5 together are a Z_2 group

$-C-R_{14}(R_{15})-C-R_{16}(R_{17})-O-C-R_{18}(R_{19})-C-R_{20}(R_{21})-$ (Z_2) wherein R_{14} , R_{15} , R_{16} , R_{17} , R_{18} , R_{19} , R_{20} and R_{21} are especially preferably hydrogen.

In a further preferred group of compositions according to the invention, in the herbicides of formula I R_{30} , R_{31} , R_{32} and R_{33} are, each independently of the others, hydrogen, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, C_1 - C_8 cyanoalkyl, C_1 - C_8 nitroalkyl, C_1 - C_8 aminoalkyl, C_2 - C_5 alkenyl, C_2 - C_5 haloalkenyl, C_3 - C_8 cycloalkyl, C_1 - C_5 alkylamino- C_1 - C_5 alkyl, C_2 - C_8 dialkylamino- C_1 - C_5 alkyl, C_3 - C_7 cycloalkyl- C_1 - C_5 alkyl, C_2 - C_4 alkoxy-alkyl, C_4 - C_6 alkenyloxy-alkyl, C_4 - C_6 alkynyloxy-alkyl, C_2 - C_4 alkylthio-alkyl, C_1 - C_4 alkylsulfinyl- C_1 - C_2 alkyl, C_1 - C_2 alkylsulfonyl- C_1 - C_2 alkyl, C_2 - C_4 alkylideneamino-oxy- C_1 - C_2 alkyl, C_1 - C_5 alkylcarbonyl- C_1 - C_2 alkyl, C_1 - C_5 alkoxycarbonyl- C_1 - C_2 alkyl, C_1 - C_5 amino-carbonyl- C_1 - C_2 alkyl, C_2 - C_8 dialkylamino-carbonyl- C_1 - C_2 alkyl, C_1 - C_5 alkylcarbonyl-amino- C_1 - C_2 alkyl, C_2 - C_5 alkylcarbonyl-(C_1 - C_2 alkyl)-aminoalkyl, C_3 - C_6 trialkylsilyl- C_1 - C_5 alkyl, phenyl- C_1 - C_2 alkyl, heteroaryl- C_1 - C_2 alkyl, phenoxy- C_1 - C_2 alkyl, heteroaryloxy- C_1 - C_2 alkyl, phenyl or heteroaryl;

R_{34} , R_{35} and R_{36} , are each independently of the others, hydrogen, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, C_1 - C_8 cyanoalkyl, C_1 - C_8 nitroalkyl, C_1 - C_8 aminoalkyl, C_2 - C_5 alkenyl, C_2 - C_5 haloalkenyl, C_3 - C_8 cycloalkyl, C_1 - C_5 alkylamino- C_1 - C_5 alkyl, C_2 - C_8 dialkylamino- C_1 - C_5 alkyl, C_3 - C_7 cycloalkyl- C_1 - C_5 alkyl, C_2 - C_4 alkoxy-alkyl, C_4 - C_6 alkenyloxy-alkyl, C_4 - C_6 alkynyloxy-alkyl, C_2 - C_4 alkylthio-alkyl, C_1 - C_4 alkylsulfinyl- C_1 - C_2 alkyl, C_1 - C_2 alkylsulfonyl- C_1 - C_2 alkyl, C_2 - C_4 alkylideneamino-oxy- C_1 - C_2 alkyl, C_1 - C_5 alkylcarbonyl- C_1 - C_2 alkyl, C_1 - C_5 alkoxycarbonyl- C_1 - C_2 alkyl, C_1 - C_5 amino-carbonyl- C_1 - C_2 alkyl, C_2 - C_8 dialkylamino-carbonyl- C_1 - C_2 alkyl, C_1 - C_5 alkylcarbonylamino- C_1 - C_2 alkyl, C_2 - C_5 alkylcarbonyl-(C_1 - C_2 alkyl)-aminoalkyl, C_3 - C_6 trialkylsilyl- C_1 - C_5 alkyl, phenyl- C_1 - C_2 -

alkyl, heteroaryl-C₁-C₂alkyl, phenoxy-C₁-C₂alkyl, heteroaryloxy-C₁-C₂alkyl, phenyl or heteroaryl, benzyloxy or phenoxy, it being possible for the benzyl and phenyl groups themselves to be substituted by halogen, nitro, cyano, amino, dimethylamino, hydroxy, methoxy, ethoxy, methylthio, ethylthio, formyl, acetyl, propionyl, carboxyl, C₁-C₅alkoxy-carbonyl or by C₁- or C₂-haloalkyl; and

R₃₇ is C₁-C₈alkyl, C₁-C₈haloalkyl, C₁-C₈cyanoalkyl, C₁-C₈nitroalkyl, C₁-C₈aminoalkyl, C₂-C₅-alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, C₁-C₅alkylamino-C₁-C₅alkyl, C₂-C₈dialkylamino-C₁-C₅alkyl, C₃-C₇cycloalkyl-C₁-C₅alkyl, C₂-C₄alkoxy-alkyl, C₄-C₆alkenyloxy-alkyl, C₄-C₆-alkynyloxy-alkyl, C₂-C₄alkylthio-alkyl, C₁-C₄alkylsulfinyl-C₁-C₂alkyl, C₁-C₂alkylsulfonyl-C₁-C₂-alkyl, C₂-C₄alkylideneamino-oxy-C₁-C₂alkyl, C₁-C₅alkylcarbonyl-C₁-C₂alkyl, C₁-C₅alkoxy-carbonyl-C₁-C₂alkyl, C₁-C₅amino-carbonyl-C₁-C₂alkyl, C₂-C₈dialkylamino-carbonyl-C₁-C₂alkyl, C₁-C₅alkylcarbonylamino-C₁-C₂alkyl, C₂-C₅alkylcarbonyl-(C₁-C₂alkyl)-aminoalkyl, C₃-C₆trialkyl-silyl-C₁-C₅alkyl, phenyl-C₁-C₂alkyl, heteroaryl-C₁-C₂alkyl, phenoxy-C₁-C₂alkyl, heteroaryloxy-C₁-C₂alkyl, phenyl or heteroaryl, benzyloxy or phenoxy, it being possible for the benzyl and phenyl groups themselves to be substituted by halogen, nitro, cyano, amino, dimethylamino, hydroxy, methoxy, ethoxy, methylthio, ethylthio, formyl, acetyl, propionyl, carboxyl, C₁-C₂-alkoxycarbonyl or by C₁- or C₂-haloalkyl; or R₃₇ is C₁-C₈alkylcarbonyl.

Special preference is given to those compositions according to the invention wherein, in the herbicides of formula I, R₃₀, R₃₁, R₃₂ and R₃₃ are, each independently of the others, hydrogen, C₁-C₈alkyl, C₁-C₈haloalkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, C₃-C₇cycloalkyl-C₁-C₂alkyl, C₂-C₄alkoxy-alkyl, phenyl, heteroaryl, phenyl-C₁-C₂alkyl, heteroaryl-C₁-C₂alkyl, phenoxy-C₁-C₂alkyl, heteroaryloxy-C₁-C₂alkyl;

R₃₄, R₃₅ and R₃₆ are, each independently of the others, hydrogen, C₁-C₈alkyl, C₁-C₈haloalkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, C₃-C₇cycloalkyl-C₁-C₂alkyl, C₂-C₄alkoxy-alkyl, phenyl, heteroaryl, phenyl-C₁-C₂alkyl, heteroaryl-C₁-C₂alkyl, phenoxy-C₁-C₂alkyl, heteroaryloxy-C₁-C₂alkyl, C₁-C₆alkoxy, C₁-C₃alkylamino or di(C₁-C₃alkyl)amino; and

R₃₇ is C₁-C₈alkyl, C₁-C₈haloalkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, C₃-C₇-cycloalkyl-C₁-C₂alkyl, C₂-C₄alkoxy-alkyl, phenyl, heteroaryl, phenyl-C₁-C₂alkyl, heteroaryl-C₁-C₂alkyl, phenoxy-C₁-C₂alkyl, heteroaryloxy-C₁-C₂alkyl, C₁-C₆alkoxy, C₁-C₃alkylamino, di-(C₁-C₃alkyl)-amino or C₁-C₈alkylcarbonyl.

A further group of especially preferred compositions comprise as herbicides of formula I those wherein R_1 and R_3 are ethyl, R_4 and R_5 together are a group Z_2 $-C-R_{14}(R_{15})-C-R_{16}(R_{17})-O-C-R_{18}(R_{19})-C-R_{20}(R_{21})-$, wherein R_{14} , R_{15} , R_{16} , R_{17} , R_{18} , R_{19} , R_{20} and R_{21} are hydrogen, and G is hydrogen or a radical of formula $-C(X_1)-R_{30}$ wherein X_1 is oxygen and R_{30} is hydrogen or C_1 - C_8 alkyl, especially C_4 alkyl, preferably tert-butyl.

The compositions according to the invention may also comprise salts that the compounds of formula I may form with acids. Suitable acids for the formation of acid addition salts are both organic and inorganic acids. Examples of such acids are hydrochloric acid, hydrobromic acid, nitric acid, phosphoric acids, sulfuric acid, acetic acid, propionic acid, butyric acid, valeric acid, oxalic acid, malonic acid, fumaric acid, organic sulfonic acids, lactic acid, tartaric acid, citric acid and salicylic acid. The salts of the compounds of formula I having acid hydrogen are also alkali metal salts, for example sodium and potassium salts; alkaline earth metal salts, for example calcium and magnesium salts; ammonium salts, that is to say unsubstituted ammonium salts and mono- or poly-substituted ammonium salts, or are salts with other organic nitrogen bases. Suitable salt formers are accordingly alkali metal and alkaline earth metal hydroxides, especially the hydroxides of lithium, sodium, potassium, magnesium or calcium, with those of sodium or potassium being given special importance.


Examples of suitable amines for ammonium salt formation that come into consideration are ammonia as well as primary, secondary and tertiary C_1 - C_{16} alkylamines, C_1 - C_4 hydroxyalkylamines and C_2 - C_4 alkoxyalkylamines, for example methylamine, ethylamine, n-propylamine, isopropylamine, the four butylamine isomers, n-amylamine, isoamylamine, hexylamine, heptylamine, octylamine, nonylamine, decylamine, pentadecylamine, hexadecylamine, heptadecylamine, octadecylamine, methyl-ethylamine, methyl-isopropylamine, methyl-hexylamine, methyl-nonylamine, methyl-pentadecylamine, methyl-octadecylamine, ethyl-butylamine, ethyl-heptylamine, ethyl-octylamine, hexyl-heptylamine, hexyl-octylamine, dimethylamine, diethylamine, di-n-propylamine, diisopropylamine, di-n-butylamine, di-n-amylamine, diisoamylamine, dihexylamine, diheptylamine, dioctylamine, ethanolamine, n-propanolamine, isopropanolamine, N,N-diethanolamine, N-ethylpropanolamine, N-butyl-ethanolamine, allylamine, n-butenyl-2-amine, n-pentenyl-2-amine, 2,3-dimethylbutenyl-2-amine, dibutenyl-2-amine, n-hexenyl-2-amine, propylenediamine, trimethylamine, triethylamine, tri-n-propylamine, triisopropylamine, tri-n-butylamine, triisobutylamine, tri-sec-

butylamine, tri-n-amyamine, methoxyethylamine and ethoxyethylamine; heterocyclic amines, for example pyridine, quinoline, isoquinoline, morpholine, N-methylmorpholine, thio-morpholine, piperidine, pyrrolidine, indoline, quinuclidine and azepine; primary aryl amines, for example anilines, methoxyanilines, ethoxyanilines, o-, m- and p-toluidines, phenylene-diamines, benzidines, naphthylamines and o-, m- and p-chloroanilines; but especially triethylamine, isopropylamine and diisopropylamine.


If non-chiral starting materials are employed, the asymmetrically substituted compounds of formula I obtained in the processes described in this Application are generally in the form of racemates. The stereoisomers can then be separated on the basis of their physicochemical properties according to known methods, such as, for example, fractional crystallisation following salt formation with optically pure bases, acids or metal complexes, or by chromatographic procedures, such as, for example, high-pressure liquid chromatography (HPLC) on acetyl cellulose. In the present invention, "compounds of formula I" are to be understood as including both the concentrated and optically pure forms of the stereoisomers in question and the racemates and diastereoisomers. Where no special mention is made of individual optical antipodes, the formula in question is to be understood as referring to the racemic mixtures that are obtained in the preparation process mentioned. When an aliphatic C=C double bond is present, geometric isomerism may also occur.

The compounds of formula I may, also in dependence upon the nature of the substituents, occur as geometric and/or optical isomers and isomeric mixtures and as tautomers and tautomeric mixtures. For example, the compounds of formula I wherein the group G is hydrogen can occur in the following tautomeric equilibria:

- 11 -




When G is other than hydrogen and Z is the group Z₁ or Z₃, or when G is other than hydrogen and Z₂ is asymmetrically substituted, fused or spiro-bound, the compound of formula I may occur as an isomer of formula Id



Processes for the preparation of compounds that differ from the compounds of formula I according to the present invention in respect of the meanings of the substituents R₄ and R₅ are described, for example, in WO 96/21652. The compounds of formula I according to the present invention can be prepared analogously to the processes described in WO 99/47525 and WO 01/17351.

Table 1: Compounds of formula Ia:

(The substituents R_4 and R_5 in the compounds of formula I form a $-C_2H_4-O-C_2H_4-$ radical in the compounds of formula Ia)






(Ia)

Comp. No.	R_1	R_3	G	Physical data
1.001	CH ₃	OCH ₃	H	
1.002	CH ₃	OCH ₃	C(O)C(CH ₃) ₃	
1.003	CH ₃	OCH ₃	C(O)OCH ₂ CH ₃	
1.004	CH ₂ CH ₃	CH ₃	H	m.p. 182-185°C
1.005	CH ₂ CH ₃	CH ₃	C(O)C(CH ₃) ₃	m.p. 110-113°C
1.006	CH ₂ CH ₃	CH ₃	C(O)OCH ₂ CH ₃	
1.007	CH ₂ CH ₃	CH ₂ CH ₃	H	m.p. 189-191°C
1.008	CH ₂ CH ₃	CH ₂ CH ₃	C(O)C(CH ₃) ₃	m.p. 122-124°C
1.009	CH ₂ CH ₃	CH ₂ CH ₃	C(O)OCH ₂ CH ₃	m.p. 114-116°C
1.010	CH=CH ₂	CH ₃	H	m.p. 165-170°C





- 13 -

Comp. No.	R ₁	R ₃	G	Physical data
1.011	CH=CH ₂	CH ₃	C(O)C(CH ₃) ₃	m.p. 111-113°C
1.012	CH=CH ₂	CH ₂ CH ₃	H	
1.013	CH=CH ₂	CH=CH ₂	H	
1.014	CH=CH ₂	CH=CH ₂	C(O)C(CH ₃) ₃	
1.015	C≡CH	CH ₃	H	m.p. 179-184°C
1.016	C≡CH	CH ₃	C(O)C(CH ₃) ₃	m.p. 109-111°C
1.017	C≡CH	CH ₃	C(O)OCH ₂ CH ₃	
1.018	C≡CH	CH ₂ CH ₃	H	m.p. 189-193°C
1.019	C≡CH	CH ₂ CH ₃	C(O)C(CH ₃) ₃	
1.020	C≡CH	CH ₂ CH ₃	C(O)OCH ₂ CH ₃	
1.021	C≡CH	C≡CH	H	m.p. 300°C
1.022	C≡CH	C≡CH	C(O)C(CH ₃) ₃	m.p. 183-185°C
1.023	C≡CH	C≡CH	C(O)OCH ₂ CH ₃	
1.024	C≡CH	CH=CH ₂	H	
1.025	C≡CCH ₃	CH ₃	H	m.p. 179-181°C
1.026	C≡CCH ₃	CH ₃	C(O)C(CH ₃) ₃	m.p. 128-129°C
1.027	C≡CCH ₃	CH ₃	C(O)OCH ₂ CH ₃	
1.028	C≡CCH ₃	CH ₂ CH ₃	H	
1.029	C≡CCH ₃	CH ₂ CH ₃	C(O)C(CH ₃) ₃	
1.030	C≡CCH ₃	C≡CCH ₃	H	
1.031	C≡CCH ₃	C≡CCH ₃	C(O)C(CH ₃) ₃	



- 14 -

Comp. No.	R ₁	R ₃	G	Physical data
1.032	CH ₂ CH ₂ CH ₃	CH ₃	H	m.p. 136-138°C
1.033	CH ₂ CH ₂ CH ₃	CH ₃	C(O)C(CH ₃) ₃	m.p. 65-67°C
1.034	CH ₂ CH ₂ CH ₃	CH ₃	C(O)OCH ₂ CH ₃	
1.035	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	H	
1.036	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	H	
1.037	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	C(O)C(CH ₃) ₃	
1.038	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	C(O)OCH ₂ CH ₃	
1.039	CH ₂ CH ₂ CH ₃	C≡CH	H	
1.040	CH(CH ₃) ₂	CH ₃	H	m.p. 214-216°C
1.041	CH(CH ₃) ₂	CH ₃	C(O)C(CH ₃) ₃	m.p. 148-151°C
1.042	CH(CH ₃) ₂	CH ₂ CH ₃	H	
1.043	CH(CH ₃) ₂	C≡CH	H	
1.044		CH ₃	H	
1.045		CH ₂ CH ₃	H	
1.046		C≡CH	H	
1.047	CH ₂ CH=CH ₂	CH ₃	H	
1.048	CH ₂ CH=CH ₂	CH ₂ CH ₃	H	
1.049	CH ₂ CH=CH ₂	C≡CH	H	
1.050	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃	H	
1.051	CH ₃ O-	CH ₂ CH ₃	H	
1.052	CH ₃ O-	CH ₂ CH ₃	C(O)C(CH ₃) ₃	
1.053	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH(CH ₃) ₂	
1.054	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH ₃	crystalline
1.055	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH(CH ₃) ₂	
1.056	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CF ₃	





- 15 -

Comp. No.	R ₁	R ₃	G	Physical data
1.057	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH ₂ CH ₃	
1.058	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH ₂ CH(CH ₃) ₂	wax
1.059	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH ₂ CH ₂ Cl	
1.060	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH=CH ₂	wax
1.061	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH ₂ CH ₂ Br	
1.062	CH ₂ CH ₃	CH ₂ CH ₃		m.p.:204-205
1.063	CH ₂ CH ₃	CH ₂ CH ₃		m.p.:203-204
1.064	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ -benzyl	m.p.:157-158
1.065	CH ₂ CH ₃	CH ₂ CH ₃		wax
1.066	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH ₂ CH ₂ CH ₂ Cl	wax
1.067	CH ₂ CH ₃	CH ₂ CH ₃		m.p.: 126

- 16 -

Comp. No.	R ₁	R ₃	G	Physical data
1.068	CH ₂ CH ₃	CH ₂ CH ₃		m.p.: 146
1.069	CH ₂ CH ₃	CH ₂ CH ₃		m.p.: 82-85
1.070	CH ₂ CH ₃	CH ₂ CH ₃	SO ₂ CH ₂ CH=CH ₂	
1.071	C≡CH	CH ₂ CH ₃	SO ₂ CH ₃	
1.072	C≡CH	CH ₂ CH ₃	SO ₂ CH(CH ₃) ₂	
1.073	C≡CH	CH ₂ CH ₃	SO ₂ CH ₂ CH ₂ Cl	
1.074	C≡CH	CH ₂ CH ₃	SO ₂ CF ₃	
1.075	C≡CH	CH ₂ CH ₃	SO ₂ CH=CH ₂	
1.076	C≡CH	OCH ₃	-H	m.p. 202-204
1.077	C≡CH	OCH ₃	C(O)C(CH ₃) ₃	m.p. 204-206
1.078	C≡CSi(CH ₃) ₃	OCH ₃	C(O)C(CH ₃) ₃	m.p. 169-171
1.079	C≡CSi(CH ₃) ₃	OCH ₃	-H	m.p. 173-174
1.080	Br	OCH ₃	-H	m.p. 217-219
1.081	Br	OCH ₃	C(O)C(CH ₃) ₃	m.p. 173-175
1.082	CH ₂ CH ₃	CH ₂ CH ₃	C(O)C(CH ₃) ₂ CH ₂ CH ₃	m.p. 122-124°C
1.083	CH ₂ CH ₃	CH ₂ CH ₃	CON(CH ₂ CH ₃) ₂	m.p. 82-84

- 17 -

Comp. No.	R ₁	R ₃	G	Physical data
1.084	CH ₂ CH ₃	C(O)CH ₃	C(O)C(CH ₃) ₂ CH ₂ CH ₃	m.p. 138-139°C
1.085	CH ₂ CH ₃	C(O)CH ₃		
1.086	CH ₂ CH ₃	C(O)CH ₃		
1.087	CH ₂ CH ₃	C(O)CH ₃		
1.088	CH ₂ CH ₃	C(O)CH ₃		

The rates of application of herbicide are generally from 0.001 to 2 kg/ha, but preferably from 0.005 to 1 kg/ha.

The ratio by weight of the compound of formula I to the second herbicide (mesosulfuron, mesotrione or flufenacet) in the composition according to the invention is preferably from 1 : 100 to 1000 : 1.

The compositions according to the invention preferably contain in addition a safener and, optionally, an oil additive. The present invention accordingly relates also to herbicidal compositions that, in addition to comprising customary inert formulation adjuvants, such as carriers, solvents and wetting agents, comprise as active ingredient a mixture of

- a) a herbicide of formula I,
- b) an amount, effective for herbicide synergy, of mesosulfuron, mesotrione or flufenacet,
- c) an amount, effective for herbicide antagonism, of a safener selected from cloquintocet-mexyl and mefenpyr-diethyl; and, optionally,
- d) an additive comprising an oil of vegetable or animal origin, a mineral oil, alkyl esters of such oils, or mixtures of such oils and oil derivatives.

The safeners cloquintocet-mexyl and mefenpyr-diethyl can also be used in the form of their alkali metal, alkaline earth metal, sulfonium or ammonium salts. Examples thereof are described, for example, in WO 02/34048. It is also possible to use hydrates of cloquintocet-mexyl, which are mentioned in WO 02/36566.

Cultivated plants that can be protected against the harmful effect of the above-mentioned herbicide mixtures by means of such safeners are especially cereals, cotton, soybeans, sugar beet, sugar cane, plantation crops, rape, maize and rice, more especially cereals. Crops are to be understood as including those that have been made tolerant to herbicides or classes of herbicides by means of conventional breeding or genetic engineering methods.

The weeds to be controlled may be either dicotyledonous or, preferably, monocotyledonous weeds, for example the monocotyledonous weeds *Avena*, *Agrostis*, *Phalaris*, *Lolium*, *Bromus*, *Alopecurus*, *Setaria*, *Digitaria*, *Brachiaria*, *Echinochloa*, *Panicum*, *Sorghum* hal./bic., *Rottboellia*, *Cyperus*, *Brachiaria*, *Echinochloa*, *Scirpus*, *Monochoria* and *Sagittaria*, and the dicotyledonous weeds *Sinapis*, *Chenopodium*, *Stellaria*, *Galium*, *Viola*, *Veronica*, *Matricaria*, *Papaver*, *Solanum*, *Abutilon*, *Sida*, *Xanthium*, *Amaranthus*, *Ipomoea* and *Chrysanthemum*.

Areas of cultivation include the areas of ground on which cultivated plants are already growing or which have already been sown with the seeds of those cultivated plants, as well as ground intended for cultivation with such cultivated plants.

Depending on the intended use, a safener according to the invention can be used to pre-treat the seed material of the cultivated plant (dressing the seed or the cuttings) or can be introduced into the soil before or after sowing. It can, however, also be applied, either alone or together with the herbicide mixture and the oil additive, after the emergence of the plants. The treatment of the plants or seed with the safener can therefore, in principle, be effected independently of the time at which the herbicide mixture is applied. The treatment of the plants can, however, also be carried out by applying herbicide, oil additive and safener simultaneously (for example in the form of a tank mixture). The rate of application of safener in relation to herbicide depends largely on the method of application. In the case of field treatment, which is effected either using a tank mixture comprising a combination of safener and herbicide mixture or by separate application of safener and herbicide mixture, the ratio of herbicides to safener is generally from 100:1 to 1:10, preferably from 20:1 to 1:1. In the case of field treatment, from 0.001 to 1.0 kg of safener/ha, preferably from 0.001 to 0.25 kg of safener/ha, is generally applied.

In the composition according to the invention, the amounts of oil additive employed are generally from 0.01 to 2 %, based on the spray mixture. The oil additive can, for example, be added to the spray tank in the desired concentration after the spray mixture has been prepared.

Preferred oil additives comprise mineral oils or an oil of vegetable origin such as, for example, rapeseed oil, olive oil or sunflower oil, alkyl esters of oils of vegetable origin such as, for example, the methyl derivatives, or an oil of animal origin such as fish oil or beef tallow.

Especially preferred oil additives comprise alkyl esters of higher fatty acids (C_8 - C_{22}), especially the methyl derivatives of C_{12} - C_{18} fatty acids, for example the methyl esters of lauric acid, palmitic acid and oleic acid. Those esters are known as methyl laurate (CAS-111-82-0), methyl palmitate (CAS-112-39-0) and methyl oleate (CAS-112-62-9).

The application and action of the oil additives can be improved by combining them with surface-active substances such as non-ionic, anionic or cationic surfactants. Examples of suitable anionic, non-ionic and cationic surfactants are listed in WO 97/34485 on pages 7 and 8.

Preferred surface-active substances are anionic surfactants of the dodecylbenzylsulfonate type, especially the calcium salts thereof, and also non-ionic surfactants of the fatty alcohol ethoxylate type. Special preference is given to ethoxylated C_{12} - C_{22} fatty alcohols having a degree of ethoxylation of from 5 to 40. Examples of commercially available preferred surfactants are the Genapol types (Clariant AG, Muttenz, Switzerland).

The concentration of the surface-active substances based on the total additive is generally from 1 to 30 % by weight.

Examples of oil additives consisting of mixtures of oils or mineral oils or derivatives thereof with surfactants are Edenor ME SU®, Emery 2231® (Henkel subsidiary company Cognis GmbH, Germany), Turbocharge® (Zeneca Agro, Stoney Creek, Ontario, Canada) or, more especially, Actipron® (BP Oil UK Limited, GB).

The addition of an organic solvent to the oil additive/surfactant mixture can, furthermore, bring about a further increase in action. Suitable solvents are, for example Solvesso® (ESSO) or Aromatic Solvent® (Exxon Corporation) types.

The concentration of those solvents can be from 10 to 80 %, by weight, of the total weight.

Such oil additives, which are also described, for example, in US-A-4 834 908, are especially preferred for the composition according to the invention. An especially preferred oil additive is known under the name MERGE®, can be obtained from the BASF Corporation and is basically described, for example, in US-A-4 834 908 in col. 5, as Example COC-1. A further oil additive that is preferred according to the invention is SCORE® (Novartis Crop Protection Canada).

In the composition according to the invention, the amounts of oil additive employed are generally from 0.01 to 2 %, based on the spray mixture. The oil additive can, for example, be added to the spray tank in the desired concentration after the spray mixture has been prepared.

The invention relates also to a method for the selective control of weeds and grasses in crops of useful plants, which method comprises treating the useful plants, seeds or cuttings thereof or the area of cultivation thereof with a herbicidal composition that comprises a mixture of

- a) a herbicidally effective amount of formula I,
- b) an amount, effective for herbicide synergy, of mesosulfuron, mesotrione or flufenacet,
- c) an amount, effective for herbicide antagonism, of a safener selected from cloquintocet-mexyl and mefenpyr-diethyl; and, optionally,
- d) an additive comprising an oil of vegetable or animal origin, a mineral oil, alkyl esters of such oils, or mixtures of such oils and oil derivatives.

The compositions according to the invention are suitable for all methods of application that are customary in agriculture, for example pre-emergence application, post-emergence application and seed dressing.

In the case of seed dressing, from 0.001 to 10 g of safener/kg of seed, preferably from 0.05 to 6 g of safener/kg of seed, are generally applied. When the safener is applied in liquid form shortly before sowing, with swelling of the seed, it is advantageous to use safener solutions that comprise the active ingredient in a concentration of from 1 to 10 000 ppm, preferably from 10 to 1000 ppm.

For application, the safeners used according to the invention or combinations of those safeners with the herbicides and, optionally, the oil additives are processed, together with the adjuvants conventionally employed in formulation technology, into formulations, for example into emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules or microcapsules.

Such formulations are described, for example, in WO 97/34485, on pages 9 to 13. The formulations are prepared in known manner, for example by intimately mixing and/or grinding the active ingredients with liquid or solid formulation adjuvants, for example solvents or solid carriers. Furthermore, surface-active compounds (surfactants) may additionally be used in the preparation of the formulations. Solvents and solid carriers suitable for that purpose are mentioned, for example, in WO 97/34485 on page 6.

Depending on the nature of the compound of formula I being formulated, suitable surface-active compounds are non-ionic, cationic and/or anionic surfactants and mixtures of surfactants having good emulsifying, dispersing and wetting properties. Examples of suitable anionic, non-ionic and cationic surfactants are listed, for example, in WO 97/34485 on pages 7 and 8. Furthermore, the surfactants customarily employed in formulation technology, which are described, *inter alia*, in "Mc Cutcheon's Detergents and Emulsifiers Annual" MC Publishing Corp., Ridgewood New Jersey, 1981, Stache, H., "Tensid-Taschenbuch", Carl Hanser Verlag, Munich/Vienna, 1981 and M. and J. Ash, "Encyclopedia of Surfactants", Vol I-III, Chemical Publishing Co., New York, 1980-81, are also suitable for preparation of the herbicidal compositions according to the invention.

The herbicidal formulations generally comprise from 0.1 to 99 % by weight, especially from 0.1 to 95 % by weight, of active ingredient mixture comprising the compound of formula I, the

second synergistically effective herbicide and, optionally, the safeners used in accordance with the invention, from 0 to 2 % by weight of the oil additive used in accordance with the invention, from 1 to 99.9 % by weight of a solid or liquid formulation adjuvant and from 0 to 25 % by weight, especially from 0.1 to 25 % by weight, of a surfactant. Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

The compositions may also comprise further additives such as stabilisers, for example vegetable oils or epoxidised vegetable oils (epoxidised coconut oil, rapeseed oil or soybean oil), antifoams, for example silicone oil, preservatives, viscosity regulators, binders, tackifiers, and also fertilisers or other active ingredients. There are various suitable methods and techniques for using the safeners or compositions comprising them for protecting cultivated plants against harmful effects of the herbicides; the following are examples:

i) Seed dressing

- a) Dressing the seeds with a wettable powder formulation of safener active ingredient by shaking in a vessel until the formulation is uniformly distributed over the seed surface (dry dressing). Approximately from 1 to 500 g of safener active ingredient according to the invention (from 4 g to 2 kg of wettable powder) are used per 100 kg of seed.
- b) Dressing the seeds with an emulsifiable concentrate of safener according to method a) (wet dressing).
- c) Dressing by immersing the seed in a liquid formulation comprising from 100 to 1000 ppm of safener for from 1 to 72 hours and, if desired, subsequently drying the seeds (immersion dressing).

Dressing the seed or treating the germinated seedlings are naturally the preferred methods of application because the treatment with the active ingredient is directed wholly at the target crop. Generally from 1 to 1000 g of antidote, preferably from 5 to 250 g of antidote, are used per 100 kg of seed, although, depending on the method employed, which also allows the addition of other active ingredients or micronutrients, amounts that exceed or fall short of the specified concentration limits may be employed (repeat dressing).

ii) Application in the form of a tank mixture

A liquid formulation of a mixture of antidote and herbicide (ratio of the one to the other from 20:1 to 1:100) is used, the rate of application of herbicide being from 0.005 to 5.0 kg per hectare. The oil additive can be added to the tank mixture in an amount of, preferably, from 0.01 to 2 % by weight. Such tank mixtures are applied before or after sowing.

iii) Application to the seed furrow

The safener is introduced into the open, sown seed furrow in the form of an emulsifiable concentrate, a wettable powder or granules. After the seed furrow has been covered, the herbicide, optionally in combination with the oil additive, is applied pre-emergence in the normal manner.

iv) Controlled release of the active ingredient

The safener is applied in solution to granulated mineral carriers or polymerised granules (urea-formaldehyde) and dried. If desired, a coating may be applied (coated granules) which enables the active ingredient to be released in metered amounts over a predetermined period of time.

Preferred formulations have especially the following compositions (% = percent by weight; 'active ingredient mixture' denotes the mixture of compound of formula I with the synergistically effective second herbicide and, optionally, with the safeners and/or oil additives according to the invention.

Emulsifiable concentrates:

active ingredient mixture:	from 1 to 90 %, preferably from 5 to 20 %
surface-active agent:	from 1 to 30 %, preferably from 10 to 20 %
liquid carrier:	from 5 to 94 %, preferably from 70 to 85 %

Dusts:

active ingredient mixture:	from 0.1 to 10 %, preferably from 0.1 to 5 %
solid carrier:	from 99.9 to 90 %, preferably from 99.9 to 99 %

Suspension concentrates:

active ingredient mixture:	from 5 to 75 %, preferably from 10 to 50 %
water:	from 94 to 24 %, preferably from 88 to 30 %
surface-active agent:	from 1 to 40 %, preferably from 2 to 30 %

Wettable powders:

active ingredient mixture:	from 0.5 to 90 %, preferably from 1 to 80 %
surface-active agent:	from 0.5 to 20 %, preferably from 1 to 15 %
solid carrier:	from 5 to 95 %, preferably from 15 to 90 %

Granules:

active ingredient mixture:	from 0.1 to 30 %, preferably from 0.1 to 15 %
solid carrier:	from 99.5 to 70 %, preferably from 97 to 85 %

The Examples that follow illustrate the invention further. They do not limit the invention.

Formulation Examples for mixtures of herbicides and, optionally, safener and oil additive

(% = percent by weight)

F1. Emulsifiable concentrates

	a)	b)	c)	d)
active ingredient mixture	5 %	10 %	25 %	50 %
calcium dodecylbenzenesulfonate	6 %	8 %	6 %	8 %
castor oil polyglycol ether	4 %	-	4 %	4 %
(36 mol of ethylene oxide)				
octylphenol polyglycol ether	-	4 %	-	2 %
(7-8 mol of ethylene oxide)				
cyclohexanone	-	-	10 %	20 %
aromatic C ₉ -C ₁₂ hydrocarbon mixture	85 %	78 %	55 %	16 %

Emulsions of any desired concentration can be prepared from such concentrates by dilution with water.

F2. Solutions

	a)	b)	c)	d)
active ingredient mixture	5 %	10 %	50 %	90 %
1-methoxy-3-(3-methoxy-propoxy)-propane	-	20 %	20 %	-
polyethylene glycol (mol. wt. 400)	20 %	10 %	-	-

- 25 -

N-methyl-2-pyrrolidone	-	-	30 %	10 %
aromatic C ₉ -C ₁₂ hydrocarbon mixture	75 %	60 %	-	-

The solutions are suitable for application in the form of micro-drops.

<u>F3. Wettable powders</u>	a)	b)	c)	d)
active ingredient mixture	5 %	25 %	50 %	80 %
sodium lignosulfonate	4 %	-	3 %	-
sodium lauryl sulfate	2 %	3 %	-	4 %
sodium diisobutyl-naphthalenesulfonate	-	6 %	5 %	6 %
octylphenol polyglycol ether (7-8 mol of ethylene oxide)	-	1 %	2 %	-
highly dispersed silicic acid	1 %	3 %	5 %	10 %
kaolin	88 %	62 %	35 %	-

The active ingredient is mixed thoroughly with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders which can be diluted with water to give suspensions of any desired concentration.

<u>F4. Coated granules</u>	a)	b)	c)
active ingredient mixture	0.1 %	5 %	15 %
highly dispersed silicic acid	0.9 %	2 %	2 %
inorganic carrier material	99.0 %	93 %	83 %

(diameter 0.1 - 1 mm)

for example CaCO₃ or SiO₂

The active ingredient is dissolved in methylene chloride, the solution is applied to the carrier by spraying, and the solvent is subsequently evaporated off *in vacuo*.

<u>F5. Coated granules</u>	a)	b)	c)
active ingredient mixture	0.1 %	5 %	15 %
polyethylene glycol (mol. wt. 200)	1.0 %	2 %	3 %
highly dispersed silicic acid	0.9 %	1 %	2 %
inorganic carrier material	98.0 %	92 %	80 %

(diameter 0.1 - 1 mm)

for example CaCO₃ or SiO₂

The finely ground active ingredient is uniformly applied, in a mixer, to the carrier material moistened with polyethylene glycol, yielding non-dusty coated granules.

- 26 -

F6. Extruder granules

	a)	b)	c)	d)
active ingredient mixture	0.1 %	3 %	5 %	15 %
sodium lignosulfonate	1.5 %	2 %	3 %	4 %
carboxymethylcellulose	1.4 %	2 %	2 %	2 %
kaolin	97.0 %	93 %	90 %	79 %

The active ingredient is mixed with the adjuvants, and the mixture is ground, moistened with water, extruded and then dried in a stream of air.

F7. Dusts

	a)	b)	c)
active ingredient mixture	0.1 %	1 %	5 %
talcum	39.9 %	49 %	35 %
kaolin	60.0 %	50 %	60 %

Ready-to-use dusts are obtained by mixing the active ingredient with the carriers and grinding the mixture in a suitable mill.

F8. Suspension concentrates

	a)	b)	c)	d)
active ingredient mixture	3 %	10 %	25 %	50 %
ethylene glycol	5 %	5 %	5 %	5 %
nonylphenol polyglycol ether (15 mol of ethylene oxide)	-	1 %	2 %	-
sodium lignosulfonate	3 %	3 %	4 %	5 %
carboxymethylcellulose	1 %	1 %	1 %	1 %
37 % aqueous formaldehyde solution	0.2 %	0.2 %	0.2 %	0.2 %
silicone oil emulsion	0.8 %	0.8 %	0.8 %	0.8 %
water	87 %	79 %	62 %	38 %

The finely ground active ingredient is intimately mixed with the adjuvants, giving a suspension concentrate from which suspensions of any desired concentration can be obtained by dilution with water.

It is often more practical to formulate the herbicides (optionally in combination with the oil additive) and the safener separately and then, shortly before application, to bring them together in the applicator in the desired mixing ratio in the form of a "tank mixture" in water. The herbicides and the safener can also be formulated separately and, shortly before application, brought together in the applicator in the desired mixing ratio in the form of a "tank mixture" in water, with the oil additive being added thereafter.

The herbicidally selective action of the compositions according to the invention is illustrated in the following Examples.


Biological Examples

Example B1: Post-emergence test:

The test plants are grown in pots under greenhouse conditions until a post-application stage. A standard soil is used as cultivation substrate. At a post-emergence stage, the herbicides, both on their own and in admixture with safeners and/or oil additives, are applied to the test plants or to cultivated plants seed-dressed with safeners. The application is carried out using an emulsion (prepared from an emulsifiable concentrate (Example F1, c)) of the test substances. The rates of application depend on the optimum concentrations ascertained under field conditions or greenhouse conditions. The tests are evaluated after from 2 to 4 weeks (100 % action = complete destruction, 0 % action = no phytotoxic action).

What is claimed is:



1. A herbicidal composition that, in addition to comprising customary inert formulation adjuvants, comprises as active ingredient a mixture of
- a) a herbicidally effective amount of a compound of formula I



wherein

R_1 and R_3 are, each independently of the other, halogen, nitro, cyano, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkyl, C_2 - C_6 haloalkenyl, C_3 - C_6 cycloalkyl, halo-substituted C_3 - C_6 cycloalkyl, C_2 - C_6 alkoxyalkyl, C_2 - C_6 alkylthioalkyl, hydroxy, mercapto, C_1 - C_6 alkoxy, C_3 - C_6 alkenyloxy, C_3 - C_6 alkynyloxy, carbonyl, carboxyl, C_1 - C_4 alkylcarbonyl, C_1 - C_4 hydroxyalkyl, C_1 - C_4 alkoxycarbonyl, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, amino, C_1 - C_4 alkyl-amino or di(C_1 - C_4 alkyl)amino;

R_4 and R_5 together are a group



wherein R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{17} , R_{18} , R_{19} , R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} , R_{27} , R_{28} and R_{29} are, each independently of the others, hydrogen, halogen, C_1 - C_4 alkyl or C_1 - C_4 haloalkyl, it being possible for an alkylene ring, which together with the carbon atoms of the groups Z_1 , Z_2 or Z_3 contains from 2 to 6 carbon atoms and which may be interrupted by oxygen, to be either fused or spiro-bound to the carbon atoms of the groups Z_1 , Z_2 or Z_3 , or that alkylene ring bridges at least one ring atom of the groups Z_1 , Z_2 or Z_3 ;

G is hydrogen, $-C(X_1)-R_{30}$, $-C(X_2)-X_3-R_{31}$, $-C(X_4)-N(R_{32})-R_{33}$, $-SO_2-R_{34}$, an alkali metal, alkaline earth metal, sulfonium or ammonium cation, or $-P(X_5)(R_{35})-R_{36}$ or $-CH_2-X_6-R_{37}$;

X_1 , X_2 , X_3 , X_4 , X_5 and X_6 are, each independently of the others, oxygen or sulfur;

R₃₀, R₃₁, R₃₂ and R₃₃ are, each independently of the others, hydrogen, C₁-C₁₀alkyl, C₁-C₁₀-haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino-C₁-C₅alkyl, C₂-C₈dialkylamino-C₁-C₅alkyl, C₃-C₇cycloalkyl-C₁-C₅alkyl, C₂-C₁₀alkoxy-alkyl, C₄-C₁₀-alkenyloxy-alkyl, C₄-C₁₀alkynyloxy-alkyl, C₂-C₁₀alkylthio-alkyl, C₁-C₅alkylsulfoxyl-C₁-C₅alkyl, C₁-C₅alkylsulfonyl-C₁-C₅alkyl, C₂-C₈alkylideneamino-oxy-C₁-C₅alkyl, C₁-C₅alkylcarbonyl-C₁-C₅alkyl, C₁-C₅alkoxycarbonyl-C₁-C₅alkyl, C₁-C₅amino-carbonyl-C₁-C₅alkyl, C₂-C₈dialkyl-amino-carbonyl-C₁-C₅alkyl, C₁-C₅alkylcarbonylamino-C₁-C₅alkyl, C₂-C₅alkylcarbonyl-(C₁-C₅alkyl)-aminoalkyl, C₃-C₆trialkylsilyl-C₁-C₅alkyl, phenyl-C₁-C₅alkyl, heteroaryl-C₁-C₅alkyl, phenoxy-C₁-C₅alkyl, heteroaryloxy-C₁-C₅alkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, phenyl, or C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl or heteroaryl or heteroaryl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃-alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted heteroaryl-amino, diheteroaryl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diheteroaryl-amino, phenyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl-amino, diphenyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diphenyl-amino, C₃-C₇cycloalkyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkyl-amino, di-C₃-C₇cycloalkyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted di-C₃-C₇cycloalkyl-amino, C₃-C₇cycloalkoxy or C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkoxy;

R₃₄, R₃₅ and R₃₆ are, each independently of the others, hydrogen, C₁-C₁₀alkyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkylamino-C₁-C₅alkyl, C₂-C₈dialkylamino-C₁-C₅alkyl, C₃-C₇cycloalkyl-C₁-C₅alkyl, C₂-C₁₀alkoxy-alkyl, C₄-C₁₀-alkenyloxy-alkyl, C₄-C₁₀alkynyloxy-alkyl, C₂-C₁₀alkylthio-alkyl, C₁-C₅alkylsulfoxyl-C₁-C₅alkyl, C₁-C₅alkylsulfonyl-C₁-C₅alkyl, C₂-C₈alkylideneamino-oxy-C₁-C₅alkyl, C₁-C₅alkylcarbonyl-C₁-C₅alkyl, C₁-C₅alkoxycarbonyl-C₁-C₅alkyl, C₁-C₅amino-carbonyl-C₁-C₅alkyl, C₂-C₈dialkyl-amino-carbonyl-C₁-C₅alkyl, C₁-C₅alkylcarbonylamino-C₁-C₅alkyl, C₂-C₅alkylcarbonyl-(C₁-C₅alkyl)-aminoalkyl, C₃-C₆trialkylsilyl-C₁-C₅alkyl, phenyl-C₁-C₅alkyl, heteroaryl-C₁-C₅alkyl, phenoxy-C₁-C₅alkyl, heteroaryloxy-C₁-C₅alkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₈cycloalkyl, phenyl, or C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl or heteroaryl or heteroaryl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃-alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted heteroaryl-amino, diheteroaryl-

amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diheteroaryl-amino, phenyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl-amino, diphenyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diphenyl-amino, C₃-C₇cycloalkyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkyl-amino, di-C₃-C₇cycloalkyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted di-C₃-C₇cycloalkyl-amino, C₃-C₇cycloalkoxy, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀haloalkoxy, C₁-C₅alkyl-amino, C₂-C₈dialkyl-amino and benzyloxy or phenoxy, it being possible for the benzyl and phenyl groups themselves to be substituted by C₁-C₃alkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, halogen, cyano, formyl, acetyl, propionyl, carboxyl, C₁-C₅alkoxy-carbonyl, methylthio, ethylthio or by nitro;

and

R₃₇ is C₁-C₁₀alkyl, C₁-C₁₀haloalkyl, C₁-C₁₀cyanoalkyl, C₁-C₁₀nitroalkyl, C₁-C₁₀aminoalkyl, C₁-C₅alkyl-amino-C₁-C₅alkyl, C₂-C₈dialkyl-amino-C₁-C₅alkyl, C₃-C₇cycloalkyl-C₁-C₅alkyl, C₂-C₁₀alkoxy-alkyl, C₄-C₁₀alkenyloxy-alkyl, C₄-C₁₀alkynyloxy-alkyl, C₂-C₁₀alkylthio-alkyl, C₁-C₅alkylsulfoxyl-C₁-C₅alkyl, C₁-C₅alkylsulfonyl-C₁-C₅alkyl, C₂-C₈alkylideneamino-oxy-C₁-C₅alkyl, C₁-C₅alkylcarbonyl-C₁-C₅alkyl, C₁-C₅alkoxycarbonyl-C₁-C₅alkyl, C₁-C₅amino-carbonyl-C₁-C₅alkyl, C₂-C₈dialkyl-amino-carbonyl-C₁-C₅alkyl, C₁-C₅alkylcarbonylamino-C₁-C₅alkyl, C₂-C₅alkylcarbonyl-(C₁-C₅alkyl)-aminoalkyl, C₃-C₆trialkylsilyl-C₁-C₅alkyl, phenyl-C₁-C₅alkyl, heteroaryl-C₁-C₅alkyl, phenoxy-C₁-C₅alkyl, heteroaryloxy-C₁-C₅alkyl, C₂-C₅alkenyl, C₂-C₅haloalkenyl, C₃-C₆cycloalkyl, phenyl, or C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted phenyl or heteroaryl, or heteroaryl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted heteroaryl-amino, diheteroaryl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diheteroaryl-amino, phenyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted diphenyl-amino, C₃-C₇cycloalkyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkyl-amino, di-C₃-C₇cycloalkyl-amino, C₁-C₃alkyl-, C₁-C₃haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted di-C₃-C₇cycloalkyl-amino, C₃-C₇cycloalkoxy, C₁-C₃alkyl-, C₁-C₃-

haloalkyl-, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, halo-, cyano- or nitro-substituted C₃-C₇cycloalkoxy or C₁-C₁₀alkylcarbonyl; or a salt or diastereoisomer of a compound of formula I, with the proviso that R₁ and R₃ are not simultaneously methyl; and

b) an amount, effective for herbicide synergy, of at least one herbicide selected from mesosulfuron, mesotrione and flufenacet.

2. A composition according to claim 1, which comprises c) an amount, effective for herbicide antagonism, of a safener selected from cloquintocet-mexyl and mefenpyr-diethyl.

3. A composition according to claim 1, which comprises d) an additive comprising an oil of vegetable or animal origin, a mineral oil, alkyl esters of such oils or mixtures of such oils and oil derivatives.

4. A method for the selective control of weeds and grasses in crops of useful plants, which comprises treating the useful plants, seeds or cuttings thereof, or the area of cultivation thereof, with a composition according to claim 1.

5. A method according to claim 4, wherein the crops of useful plants are cereals.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number
WO 03/028466 A3

(51) International Patent Classification⁷: **A01N 43/90**,
25/32 // (A01N 43/90, 47:36, 43:824, 41:10)

(21) International Application Number: **PCT/EP02/10829**

(22) International Filing Date:
26 September 2002 (26.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1781/01 27 September 2001 (27.09.2001) CH

(71) Applicant (for all designated States except US): **SYNGENTA PARTICIPATIONS AG** [CH/CH]; Schwarzwaldallee 215, 4058 Basel (CH).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **HOFER, Urs** [CH/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, 4058 Basel (CH).

(74) Agent: **BASTIAN, Werner**; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:
30 October 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **HERBICIDAL COMPOSITION**



(57) Abstract: Herbicidal composition comprising: a) a herbicidally effective amount of a compound of formula (I), wherein the substituents are as defined in claim 1; and b) an amount, effective for herbicide synergy, of mesosulfuron, mesotrione or flufenacet.

WO 03/028466 A3

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 02/10829

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A01N43/90 A01N25/32 //(A01N43/90,47:36,43:824,41:10)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 17351 A (GLOCK JUTTA ;CORNES DEREK (CH); FRIEDMANN ADRIAN ALBERTO (CH); SYN) 15 March 2001 (2001-03-15) cited in the application page 1, paragraph 1 page 2, paragraph 2 -page 6, paragraph 1 page 10, paragraph 3 -page 12, paragraph 2 page 11, line 17	1-5
A	WO 01 17352 A (GLOCK JUTTA ;CORNES DEREK (CH); FRIEDMANN ADRIAN ALBERTO (CH); SYN) 15 March 2001 (2001-03-15) page 1, paragraph 1 page 2, paragraph 3 -page 6, paragraph 2 -/-	1-5

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the International search

16 April 2003

Date of mailing of the international search report

18. 05. 2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Lamers, W

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 02/10829

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 24320 A (BIERINGER HERMANN ;HACKER ERWIN (DE); LORENZ KLAUS (DE); HOECHST S) 11 June 1998 (1998-06-11) page 4, line 4 -page 5, line 28 page 9, line 28 -page 35, line 11 -----	1-5
A	WO 00 03597 A (HOECHST SCHERING AGREVO GMBH) 27 January 2000 (2000-01-27) page 1, line 4 - line 20 page 8, line 20 -page 60, line 2 -----	1-5
A	FR 2 781 983 A (SUMITOMO CHEMICAL CO) 11 February 2000 (2000-02-11) page 1, line 15 -page 2, line 13 page 4, line 22 -page 5, line 35 -----	1-5
A	US 6 001 774 A (MITO NOBUAKI) 14 December 1999 (1999-12-14) column 1, line 16 -column 2, line 16 -----	1-5

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 02/10829

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-5 (partially)

A herbicidal composition that, in addition to comprising customary inert formulation adjuvants, comprises as active ingredient a mixture of a) a herbicidally effective amount of a compound of formula I and b) an amount, effective for herbicide synergy, of mesosulfuron;
- compositions as described in claims 2 and 3, and methods as described in claims 4 and 5.

2. Claims: 1-5 (partially)

A herbicidal composition that, in addition to comprising customary inert formulation adjuvants, comprises as active ingredient a mixture of a) a herbicidally effective amount of a compound of formula I and b) an amount, effective for herbicide synergy, of mesotrione;
- compositions as described in claims 2 and 3, and methods as described in claims 4 and 5.

3. Claims: 1-5 (partially)

A herbicidal composition that, in addition to comprising customary inert formulation adjuvants, comprises as active ingredient a mixture of a) a herbicidally effective amount of a compound of formula I and b) an amount, effective for herbicide synergy, of flufenacet;
- compositions as described in claims 2 and 3, and methods as described in claims 4 and 5.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/10829

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0117351	A	15-03-2001	AU 7284400 A	10-04-2001
			CN 1376032 T	23-10-2002
			WO 0117351 A1	15-03-2001
			EP 1209972 A1	05-06-2002
			HU 0202828 A2	28-01-2003
WO 0117352	A	15-03-2001	AU 7417200 A	10-04-2001
			CN 1371244 T	25-09-2002
			WO 0117352 A1	15-03-2001
			EP 1209973 A1	05-06-2002
WO 9824320	A	11-06-1998	DE 19650955 A1	10-06-1998
			AU 731739 B2	05-04-2001
			AU 5483298 A	29-06-1998
			BG 103511 A	30-06-2000
			BR 9714380 A	21-03-2000
			CN 1239867 A	29-12-1999
			CZ 9902031 A3	17-11-1999
			WO 9824320 A1	11-06-1998
			EP 0946100 A1	06-10-1999
			HU 0000340 A2	28-06-2000
			NZ 335869 A	27-04-2001
			PL 333945 A1	31-01-2000
			SK 75899 A3	10-12-1999
			TR 9901272 T2	23-08-1999
			US 6221809 B1	24-04-2001
			ZA 9710859 A	08-06-1998
WO 0003597	A	27-01-2000	DE 19832017 A1	27-01-2000
			AU 4782099 A	07-02-2000
			AU 5158799 A	07-02-2000
			AU 5281299 A	07-02-2000
			BG 105138 A	31-10-2001
			BR 9912823 A	02-05-2001
			BR 9912828 A	02-05-2001
			BR 9912829 A	02-05-2001
			CA 2337521 A1	27-01-2000
			CN 1318978 T	24-10-2001
			CN 1309533 T	22-08-2001
			CN 1314785 T	26-09-2001
			WO 0003591 A2	27-01-2000
			WO 0003597 A1	27-01-2000
			WO 0003592 A2	27-01-2000
			EP 1104232 A2	06-06-2001
			EP 1096854 A2	09-05-2001
			EP 1096855 A1	09-05-2001
			HU 0104456 A2	28-03-2002
			JP 2002520340 T	09-07-2002
			JP 2002520341 T	09-07-2002
			JP 2002520342 T	09-07-2002
			PL 345665 A1	02-01-2002
			SK 522001 A3	11-09-2001
			TR 200100099 T2	21-06-2001
			TR 200100100 T2	23-07-2001
			TR 200100101 T2	21-06-2001
			US 2002123428 A1	05-09-2002
			US 6492301 B1	10-12-2002

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 02/10829

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
FR 2781983	A	11-02-2000	JP	2000053509 A		22-02-2000
			FR	2781983 A1		11-02-2000
US 6001774	A	14-12-1999	JP	10306006 A		17-11-1998
			JP	10306007 A		17-11-1998
			BR	9801579 A		11-01-2000
			CN	1200877 A		09-12-1998